

BUFFERED PEPTONE WATER pH 7.2

Dehydrated culture medium

1 - INTENDED USE

Buffered Peptone Water pH 7.2 is used as non-selective pre-enrichment medium for the procedure of detection and enumeration of Salmonella, according to FDA-BAM in foods.

2 - COMPOSITION - TYPICAL FORMULA *

(AFTER RECONSTITUTION WITH 1 L OF WATER)

Peptone 10.0 g Sodium chloride 5.0 g Disodium hydrogen phosphate anhydrous 3.5 g ^ Monopotassium phosphate 1.5 g

3 - PRINCIPLE OF THE METHOD AND EXPLANATION OF THE PROCEDURE

Buffered Peptone Water pH 7.2 is manufactured with a selected peptone, particularly rich in nutrients and with a buffer system that allows optimal recovery of microorganisms even when present in the sample in a very low number or sub-lethally injured.

The medium is prepared according to FDA-BAM medium 192 specification whose pH is set at 7.2 ± 0.2^{1} while ISO Standards (e.g ISO 6579) require a medium with pH 7.0 ± 0.2

Buffered Peptone Water pH 7.2 can be used as a non-selective pre-enrichment medium according to FDA-BAM chapter 5 for the detection of *Salmonella* in some foods and animal feeding stuffs.²

The peptone provides carbon, nitrogen, vitamins and minerals for microbial growth, sodium chloride maintains the osmotic balance, while phosphates buffer the medium at pH 7.2.

4 - DIRECTIONS FOR DEHYDRATED MEDIUM PREPARATION

Suspend 20 g in 1000 ml of cold purified water. Stir and heat if necessary to completely dissolve the powder. Distribute into flasks or tubes of suitable capacity and sterilise in the autoclave at 121°C for 15 minutes.

5 - PHYSICAL CHARACTERISTICS

Dehydrated medium appearance beige, fine, homogeneous, free-flowing powder

Prepared medium appearance pale yellow, limpid

Final pH at 20-25 °C 7.2 ± 0.2

6 - MATERIALS PROVIDED - PACKAGING

Product	Type	REF	Pack		
Buffered Peptone Water pH 7.2	Dehydrated medium	401278S2	500 g (25 L)		
	-	401278S4	5 kg (250 L)		

7 - MATERIALS REQUIRED BUT NOT PROVIDED

Autoclave, sterile loops and pipettes, incubator and laboratory equipment as required, Erlenmeyer flasks, tubes, bottles, ancillary culture media and reagents.

8 - SPECIMENS

Foods, animal feeding stuffs. Refer to applicable international standard² for the collection, transport, storage of samples and operate in accordance with good laboratory practice.

9 - TEST PROCEDURE

For details of sample preparation and enrichment, please refer to the Standard cited according to the intended use.²

For the pre-enrichment for Salmonella, in general, 225 mL of Buffered Peptone Water pH 7.2 are inoculated with 25 g of the sample, then incubated for 24 ± 2 h at 35°C.

10 - READING AND INTERPRETATION

Microbial growth in Buffered Peptone Water pH 7.2 is evidenced by the development of turbidity in the broth.

11 - USER QUALITY CONTROL

All manufactured lots of the product are released for sale after the Quality Control has been performed to check the compliance with the specifications. However, the end user can perform its own Quality Control in accordance with the local applicable regulations, in compliance with accreditation requirements and the experience of the Laboratory. The choice of strains depends on the intended use. Consult the Standard cited.¹

12 - PERFORMANCES CHARACTERISTICS

Prior to release for sale a representative sample of all lots of dehydrated and ready to use Buffered Peptone Water pH 7.2 (Test Batch: TB), is tested for productivity by comparing the results with a previously approved Reference Batch (RB).

Productivity is tested by dilution to extinction method, by inoculating 1 mL of appropriate decimal dilutions of organisms in test tubes and incubating at 35-37° for 18-24 hours and recording the highest dilution showing growth in Reference Batch (GrRB) and in Test Batch (GrTB). Productivity is tested with the following strains: S.Typhimurium ATCC 14028, S.Enteritidis ATCC 13076, S.arizonae ATCC 13314, S.Dublin wild strain CB 9.2. The productivity index GrRB-GrTB for each test strain shall be ≤ 1. CB: Biolife culture collection.

^{*}The formula may be adjusted and/or supplemented to meet the required performances criteria.

 $^{^{\}mbox{\sc h}}$ equivalent to disodium hydrogen phosphate dodecahydrate 9 g/L

Instructions for use

TS-401278S rev 2 2022/06 page 2 / 2



13 - LIMITATIONS OF THE METHOD

- Buffered Peptone Water pH 7.2 is a general-purpose medium without selective properties. Suitable selective liquid and solid media must be inoculated with the growth obtained in Buffered Peptone Water pH 7.2.
- Th test sample may increase the turbidity of the medium although bacterial growth is not present. Subculture to appropriate media is necessary to verify growth of organisms.

14 - PRECAUTIONS AND WARNINGS

- This product is for microbiological control and for professional use only; it is to be used by adequately trained and qualified laboratory personnel, observing approved biohazard precautions and aseptic techniques.
- Dehydrated media must be handled with suitable protection. Before use, consult the Safety Data Sheet.
- This culture medium contains raw materials of animal origin. The ante and post mortem controls of the animals and those during the production and distribution cycle of the raw materials, cannot completely guarantee that the product doesn't contain any transmissible pathogen. Therefore, it is recommended that the culture medium be treated as potentially infectious, and handled observing the usual specific precautions: do not ingest, inhale, or allow to come into contact with skin, eyes, mucous membranes. Download the TSE Statement from the website www.biolifeitaliana.it, describing the measures implemented by Biolife Italiana for the risk reduction linked to infectious animal diseases.
- Apply Good Manufacturing Practice in the production process of prepared media.
- All laboratory specimens should be considered infectious.
- The laboratory area must be controlled to avoid contaminants such as culture medium or microbial agents.
- Sterilize all biohazard waste before disposal. Dispose the unused medium and the sterilized plates inoculated with samples or microbial strains in accordance with current local legislation.
- Do not use the culture medium as active ingredient for pharmaceutical preparations or as production material intended for human and animal consumption.
- The Certificates of Analysis and the Safety Data Sheet of the product are available on the website www.biolifeitaliana.it.
- The information provided in this document has been defined to the best of our knowledge and ability and represents a guideline for the proper use of the product but without obligation or liability. In all cases existing local laws, regulations and standard procedures must be observed for the examination of samples collected from human and animal organic districts, for environmental samples and for products intended for human or animal consumption. Our information does not relieve our customers from their responsibility for checking the suitability of our product for the intended purpose.

15 - STORAGE CONDITIONS AND SHELF LIFE

Upon receipt, store at +10°C /+30°C away from direct light in a dry place. If properly stored, it may be used up to the expiration date. Do not use beyond this date. Avoid opening the bottle in humid places. After use, the container must be tightly closed. Discard the product if the container and/or the cap are damaged, or if the container is not well closed, or in case of evident deterioration of the powder (colour changes, hardening, large lumps).

The user is responsible for the manufacturing and quality control processes of prepared media and the validation of their shelf life, according to the type (tubes/flasks) and the applied storage conditions (temperature and packaging). According to ISO 6579-1² autoclaved Buffered Peptone Water may be stored in closed containers at 2-8 °C for up to six months.

16 - REFERENCES

- 1. FDA BAM Media M192: Buffered Peptone Water (BPW) Content current as of: 03/27/2018
- 2. FDA-BAM. Chapter 5: Salmonella. Content current as of: 03/18/2022
- 3. ISO 6579-1:2017 Microbiology of the food chain -- Horizontal method for the detection, enumeration and serotyping of Salmonella -- Part 1: Detection of Salmonella spp.

TABLE OF APPLICABLE SYMBOLS

REF Catalo	or REF	LOT	Batch code	***	Manufacturer	*	Store in a dry place	\sum	Use by
	Temperature limitation	Σ	Contents sufficient for <n> tests</n>		Consult Instructions for Use	淡	Keep away from direct light		

REVISION HISTORY

Version	Description of changes	Date	
Revision 2	Updated layout and content	2022/06	

Note: minor typographical, grammatical, and formatting changes are not included in the revision history