

TRIPLE SUGAR IRON AGAR ISO

Dehydrated and ready-to-use culture medium

1 - INTENDED USE

For the differentiation of *Enterobacteriaceae*, especially *Salmonella*, based on carbohydrate fermentation and production of hydrogen sulphide.

2 - COMPOSITION - TYPICAL FORMULA*

(AFTER RECONSTITUTION WITH 1 L OF WATER)

Beef extract	3.0 g
Yeast extract	3.0 g
Peptone	20.0 g
Lactose	10.0 g
Sucrose	10.0 g
Glucose	1.0 g
Iron (III) ammonium citrate	0.3 g
Sodium chloride	5.0 g
Sodium thiosulphate	0.3 g
Agar	12.5 g
Phenol red	24.00 mg

*The formula may be adjusted and/or supplemented to meet the required performances criteria.

3 - PRINCIPLE OF THE METHOD AND EXPLANATION OF THE PROCEDURE

The formulation of Triple Sugar Iron Agar medium is based on several microbiologists' attempts to develop a medium to aid in the identification of intestinal Gram-negative bacilli: Russel¹, Kligler², Krunweide and Kohn³. In 1940, Sulkin and Willet⁴ modified the triple sugar medium of Krunweide and Kohn by the addition of H₂S indicators. The current formulation of triple sugar iron medium is essentially a modification of Haja⁵ to Sulkin and Willet triple sugar ferrous sulphate medium.

Triple Sugar Iron (TSI) Agar ISO is intended for the differentiation of *Enterobacteriaceae*, especially *Salmonella* spp., grown on primary isolation media, based on the fermentation of glucose, lactose and sucrose, with production of acids and gas, and the production of hydrogen sulphide.^{6,7}

The culture medium is prepared according to the formula included in ISO 6579⁷ and differs from the classical TSI formulation (REF 402141) by the additional presence of beef extract and yeast extract and slightly different concentrations of sodium thiosulphate and phenol red.

The fermentation of the three carbohydrates can take place both on the surface of the slant and in the butt with or without the presence of gas (CO₂ + H₂) and 3 reaction models can be registered:

1-fermentation of glucose; 2-fermentation of glucose, lactose and/or sucrose; 3-no fermentation.⁸

In the first case, after 18-24 hours of incubation, an alkaline reaction on the slant and an acid reaction in the butt is observed. The complete consumption of glucose, present at a concentration of 0.1%, on the surface, where aerobic conditions exist, after 18-24 hours induces the oxidative degradation of peptones, with production of ammonia, alkalinity and a red colour change of phenol red (reversal of the acid-alkaline reaction). However, in the anaerobic butt the bacteria metabolize the glucose producing ATP and pyruvate, which is converted into stable acid end-products with a colour change of the indicator to yellow (acid pH).

In the second case, the microorganisms ferment glucose and one or both lactose and sucrose: after 18-24 hours of incubation an acid reaction is recorded on the slant and in the butt. This is due to the high concentration of lactose and sucrose: after 18-24 hours their degradation is not exhausted on the surface and therefore there is no utilisation of peptones and therefore no reversal of the reaction.

In the third model an alkaline reaction is recorded both on the slant and in the butt. This behaviour is not typical of *Enterobacteriaceae* but of some non-enteric non fermenting Gram-negative bacteria that can utilise the peptones for growing (*Alcaligenes faecalis*, *Acinetobacter*, *Pseudomonas*). If the degradation of the peptones is anaerobic the indicator will turn to red (alkaline pH) both on the surface and in the butt, if the degradation is aerobic, there is no colour change of phenol red in the butt.

Ferrous ammonium sulphate is an indicator of the formation of hydrogen sulphide. Thiosulphate reductase producing organisms cause the release of a sulphide molecule from the sodium thiosulfate. The hydrogen sulphide will react with ferric ions in the medium to produce iron sulphide, a black insoluble precipitate.

4- DIRECTIONS FOR MEDIUM PREPARATION

Suspend 65 g in 1000 mL of cold purified water. Heat to boiling with frequent agitation to dissolve completely. Distribute into tubes and sterilize by autoclaving at 121°C for 15 minutes. Cool in a slanted position to obtain deep butts and short slopes.

5 - PHYSICAL CHARACTERISTICS

Dehydrated medium appearance	pinkish, fine, homogeneous, free-flowing powder
Solution and prepared tubes appearance	red-orange, limpid
Final pH at 20-25 °C	7.4 ± 0.2

6 - MATERIALS PROVIDED - PACKAGING

Product	Type	REF	Pack
Triple Sugar Iron Agar ISO	Dehydrated culture medium	402141S2	500 g (7.7 L)
Triple Sugar Iron Agar ISO	Ready-to-use tubes	552141S	20 slanted tubes

7 - MATERIALS REQUIRED BUT NOT PROVIDED

Autoclave and water-bath, sterile needles, screw capped tubes, incubator and laboratory equipment as required, ancillary culture media and reagents for complete identification of the culture.

8 - SPECIMENS

Pure colonies from a culture on solid media.





9 - TEST PROCEDURE

With an inoculating needle, pick the centre of a well-isolated pure colony, inoculate the slant by first stabbing the butt to the bottom; withdraw the needle, and then streak the surface of the slant. Loosen the cap of the tube before incubating. Incubate aerobically at 37°C for 24 h ± 3 hours.

10 - READING AND INTERPRETATION

Interpret the changes in the medium as follows:

a) butt

- yellow: glucose positive (glucose fermentation);
- red or unchanged: glucose negative (no fermentation of glucose);
- black: formation of hydrogen sulphide;
- bubbles or cracks: gas formation from glucose;

b) slant surface

- yellow: lactose and/or sucrose positive (lactose and/or sucrose fermentation);
- red or unchanged: lactose and sucrose negative (no fermentation of lactose or sucrose).

The majority of the typical *Salmonella* cultures show alkaline (red) slants and acid (yellow) butts with gas formation (bubbles) and (in about 90 % of the cases) formation of hydrogen sulphide (blackening of the agar).

When a lactose-positive *Salmonella* is isolated, the TSI slant is yellow.

Interpretation of TSI reactions with *Salmonella* spp.⁷

	Acid from glucose	Gas from glucose	Acid from lactose	Acid from sucrose	H ₂ S production
S.Typhi	+	-	-	-	-
S.Paratyphi A	+	+	-	-	-
S.Paratyphi B	+	+	-	-	+
S.Paratyphi C	+	+	-	-	+
S.Gallinarum biovar gallinarum	+	+	-	-	V
S.Gallinarum biovar pullorum	+	+	-	-	V
Other <i>Salmonella</i> spp.	+	+	-	-	+

Notes

Not all isolates of *Salmonella* serovars show the reactions marked + or -. Reactions may also vary between and within serovars.

Salmonella Typhi is anaerogenic.

V = Variable results.

11 - USER QUALITY CONTROL

All manufactured lots of the product are released for sale after the Quality Control has been performed to check the compliance with the specifications. However, the end user can perform its own Quality Control in accordance with the local applicable regulations, in compliance with accreditation requirements and the experience of the Laboratory. Here below are listed some test strains useful for the quality control.

S. Typhimurium ATCC 14028: growth, red slant, yellow butt, gas +, H₂S +
S. flexneri ATCC 12022: growth, red slant, yellow butt, gas -, H₂S -
P. aeruginosa ATCC 10145: growth, red or unchanged slant, red or unchanged butt, gas -, H₂S -
Aerobic incubation at 37°C for 24 h.

ATCC is a trademark of American Type Culture Collection

12 - PERFORMANCES CHARACTERISTICS

Prior to release for sale, a representative sample of all lots of dehydrated Triple Sugar iron Agar ISO is tested for performances characteristics comparing the results with a previously approved Reference Batch.

Pure colonies cultivated on Tryptic Soy Agar of 7 *Enterobacteriaceae* strains are inoculated into the tubes: S. Enteritidis ATCC 13076, S. Typhimurium ATCC 14028, E. coli ATCC 25922, C. freundii ATCC 8090, P. rettgeri ATCC 39944, S. flexneri ATCC 12022, S. sonnei ATCC 9290, P. aeruginosa ATCC 10145. After aerobic incubation at 37°C for 24 hours, the colour changes on the slant and in the butt, the gas and H₂S production are observed. All strains show reactivity according to the specifications.

13 - LIMITATIONS OF THE METHOD

- It is necessary to inoculate the medium with a microbiological needle without breaking the agar (do not use loops).
- Perform the reading between 18 and 24 hours of incubation; early readings can induce false acidity results of the A/A type or there is not enough time for the sugar fermentation with consequent colour change of the indicator; delayed readings can give false K/K results due to the use of peptones and alkaline change of the medium.⁹
- H₂S production can mask the acid reaction in the butt, however the production of H₂S requires acidic conditions therefore the butt must be considered acid when there is blackening.
- Hydrogen sulphide production may be evident on KIA but negative on TSI. Studies by Bulmash and Fulton¹⁰ showed that the utilization of sucrose could suppress the enzymatic mechanisms responsible for H₂S production. Padron and Dockstader¹¹ found that not all H₂S-positive *Salmonella* are positive on TSI.
- An H₂S producing organism may exhibit blackening on SIM medium (positive) but none on TSI medium.⁹
- The medium does not contain inhibitors therefore a large variety of microorganisms can grow on it; for this reason, before inoculation, make sure that the organisms are catalase positive, Gram-negative bacilli.





- The addition of sucrose allows the earlier detection of coliform bacteria that ferment sucrose more rapidly than lactose. Adding sucrose also aids the identification of certain Gram-negative bacteria that could ferment sucrose but not lactose.⁸
- A pure culture is essential when inoculating the medium. If the culture is not pure, irregular results may be obtained.
- Some organisms such as the *Klebsiella-Enterobacter* group produce such an abundance of gas that the medium may be completely displaced by gas resulting in the medium being blown up into the cap. If this occurs, handle the culture with caution when sub-culturing to avoid contaminations.
- Make sure that the caps are loosened during incubation since for a correct medium performance a free exchange of air is necessary. If the caps are too closed, an acid reaction occurs only on the slant even in the presence of glucose fermentation.⁹
- Preliminary confirmation of *Salmonella* cultures shall not be based on the results of the TSI agar test only; further suitable tests are needed for a complete identification of the colonies.

14 - PRECAUTIONS AND WARNINGS

- This culture medium is for microbiological control and for professional use only; it is to be used by adequately trained and qualified laboratory personnel, observing approved biohazard precautions and aseptic techniques.
- Dehydrated media must be handled with suitable protection. Before use, consult the Safety Data Sheet.
- This culture medium contains raw materials of animal origin. The *ante* and *post mortem* controls of the animals and those during the production and distribution cycle of the raw materials, cannot completely guarantee that this product doesn't contain any transmissible pathogen. Therefore, it is recommended that the culture medium be treated as potentially infectious, and handled observing the usual specific precautions: do not ingest, inhale, or allow to come into contact with skin, eyes, mucous membranes. Download the TSE Statement from the website www.biolifeitaliana.it, describing the measures implemented by Biolife Italiana for the risk reduction linked to infectious animal diseases.
- Apply Good Manufacturing Practice in the production process of prepared media.
- Be careful when opening screw cap tubes to prevent injury due to breakage of glass.
- Ready-to-use tubes are subject to terminal sterilization by autoclaving.
- Each ready-to-use tube of this culture medium is for single use only.
- All laboratory specimens should be considered infectious.
- The laboratory area must be controlled to avoid contaminants such as medium powder or microbial agents.
- Sterilize all biohazard waste before disposal. Dispose the unused medium and the sterilized medium inoculated with samples or microbial strains in accordance with current local legislation.
- Do not use the culture medium as active ingredient for pharmaceutical preparations or as production material intended for human and animal consumption
- The Certificates of Analysis and the Safety Data Sheets of the products are available on the website www.biolifeitaliana.it.
- The information provided in this document has been defined to the best of our knowledge and ability and represents a guideline for the proper use of the product but without obligation or liability. In all cases existing local laws, regulations and standard procedures must be observed for the examination of samples collected from human and animal organic districts, for environmental samples and for products intended for human or animal consumption. Our information does not relieve our customers from their responsibility for checking the suitability of our product for the intended purpose.

15 - STORAGE CONDITIONS AND SHELF LIFE

Ready-to-use medium in tubes

Upon receipt, store tubes in their original pack at 2-8°C away from direct light. If properly stored, the tubes may be used up to the expiration date. Do not use the tubes beyond this date. Tubes from opened secondary packages can be used up to the expiration date. Opened tubes must be used immediately. Before use, check the closing and the integrity of the screw cap. Do not use tubes with signs of deterioration (e.g., microbial contamination, abnormal turbidity, precipitate, atypical colour).

Dehydrated medium

Upon receipt, store at +10°C /+30°C away from direct light in a dry place. If properly stored, it may be used up to the expiration date. Do not use beyond this date. Avoid opening the bottle in humid places. After use, the container must be tightly closed. Discard the product if the container and/or the cap are damaged, or if the container is not well closed, or in case of evident deterioration of the powder (colour changes, hardening, large lumps).

The user is responsible for the manufacturing and quality control processes of prepared media and the validation of their shelf life, according to the type (tubes/bottles) and the applied storage conditions (temperature and packaging). According to ISO 6579, the self-prepared tubes can be stored at +2°C +8°C for up to 4 weeks.⁴














16 - REFERENCES

1. Russell FF. The isolation of typhoid bacilli from urine and feces with the description of a new double sugar tube medium. J Med Res 1911; 25:21
2. Kliger IJ. A simple medium for the differentiation of members of the typhoid-paratyphoid group. Am J Public Health 1917; 7:1042-1044
3. Krumwiede C, Kohn L. A triple sugar modification of the Russell Double Sugar medium. J Med Res 1917; 37:225.
4. Sulkin SE, Willet JC. A triple sugar ferrous sulphate medium for use in identification of enteric organisms. J Lab Clin Med 1940; 25:649.
5. Hajna AA. Triple sugar iron agar medium for the identification of intestinal group of bacteria. J Bacteriol 1945; 49:516.
6. Atlas R, Parks LC. Handbook of Microbiological Media. 2nd edition CRC Press, 1997
7. ISO 6579:2017 Microbiology of the food chain — Horizontal method for the detection, enumeration and serotyping of Salmonella — Part 1: Detection of Salmonella spp.
8. Lehman D. Triple sugar iron agar protocols. 30 September 2005. American Society for Microbiology © 2016.
9. MacFaddin JF. Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria. Baltimore: Williams & Wilkins; 1985.
10. Bulmash JM, Fulton MD. Discrepant tests for hydrogen sulfide. J Bacteriol 1964; 88(2):1813
11. Padron AP, Dockstader WB. Selective medium for hydrogen sulfide production Appl Microbiol 1972; 23:1107





TABLE OF APPLICABLE SYMBOLS

 or  Catalogue number	 Batch code	 Manufacturer	 This side up	 Store in a dry place	 Fragile
 Temperature limitation	 Content sufficient for <n> tests	 Consult Instructions for Use	 Use by	 Keep away from direct light	 For single use only

REVISION HISTORY

Version	Description of changes	Date
Revision 1	Updated layout and content	2020/09

Note: minor typographical, grammatical, and formatting changes are not included in the revision history.

