

LISTERIA UVM1 ANTIMICROBIC SUPPLEMENT

LISTERIA UVM2 ANTIMICROBIC SUPPLEMENT

VIAL CONTENTS

Listeria UVM1 Antimicrobial Supplement (vial contents for 500 ml of medium)

Acriflavin 6 mg
Nalidixic Acid 10 mg

Listeria UVM2 Antimicrobial Supplement (vial contents for 500 ml of medium)

Acriflavin 12.5 mg
Nalidixic Acid 10.0 mg

DIRECTIONS

UVM1 Broth – UVM2 Broth: dissolve the contents of one vial of Listeria UVM 1 Antimicrobial Supplement or the contents of one vial of Listeria UVM 2 Antimicrobial Supplement with 5 ml of ethyl alcohol/sterile distilled water (1:1), under aseptic conditions. Add to 500 ml of Listeria UVM Enrichment Broth Base (cat. N° 401595), autoclaved and cooled to 50°C. Mix well and distribute into sterile tubes or flasks.

INTENDED USE

The selective supplements described above are used for the preparation of enrichment broths for the isolation of *Listeria* spp. from food or environmental specimens. For the details of the procedure see the technical sheet of Listeria UVM Enrichment Broth Base (cat. N° 401595).

STORAGE

Store at 2-8° - When stored as directed the supplements remains stable until the expiry date shown on the label. Do not use beyond stated expiry date.

PRECAUTIONS AND WARNING

The supplements should be used only by adequately trained personnel with knowledge of microbiological techniques in the laboratory.

PACKAGING

4240036 Listeria UVM1 Antimicrobial Supplement, 10 vials, each for 500 ml of medium
4240037 Listeria UVM2 Antimicrobial Supplement, 10vials, each for 500 ml of medium