

## **LISTERIA UVM1 ANTIMICROBIC SUPPLEMENT**

## **LISTERIA UVM2 ANTIMICROBIC SUPPLEMENT**

### **VIAL CONTENTS**

#### **Listeria UVM1 Antimicrobial Supplement** (vial contents for 500 ml of medium)

Acriflavin 6 mg  
Nalidixic Acid 10 mg

#### **Listeria UVM2 Antimicrobial Supplement** (vial contents for 500 ml of medium)

Acriflavin 12.5 mg  
Nalidixic Acid 10.0 mg

### **DIRECTIONS**

*UVM1 Broth – UVM2 Broth:* dissolve the contents of one vial of Listeria UVM 1 Antimicrobial Supplement or the contents of one vial of Listeria UVM 2 Antimicrobial Supplement with 5 ml of ethyl alcohol/sterile distilled water (1:1), under aseptic conditions. Add to 500 ml of Listeria UVM Enrichment Broth Base (cat. N° 401595), autoclaved and cooled to 50°C. Mix well and distribute into sterile tubes or flasks.

### **INTENDED USE**

The selective supplements described above are used for the preparation of enrichment broths for the isolation of *Listeria* spp. from food or environmental specimens. For the details of the procedure see the technical sheet of of Listeria UVM Enrichment Broth Base (cat. N° 401595).

### **STORAGE**

Store at 2-8° - When stored as directed the supplements remains stable until the expiry date shown on the label. Do not use beyond stated expiry date.

### **PRECAUTIONS AND WARNING**

The supplements should be used only by adequately trained personnel with knowledge of microbiological techniques in the laboratory.

### **PACKAGING**

**4240036 Listeria UVM1 Antimicrobial Supplement, 10 vials, each for 500 ml of medium**  
**4240037 Listeria UVM2 Antimicrobial Supplement, 10vials, each for 500 ml of medium**